Panacombo Dual Action Pain Relief 200mg/500mg Film-Coated Tablets

Uses

For the temporary relief of mild to moderate pain associated with migraine, headache, backache, period pain, dental pain, rheumatic and muscular pain, cold and flu symptoms, sore throat and fever. Especially suitable for pain that requires stronger analgesia than ibuprofen or paracetamol alone. Before this medicine is taken, the patient should first try ibuprofen or paracetamol for pain relief for the first day of treatment. If the pain has not been relieved by ibuprofen or paracetamol during the first day of treatment, this medicine can be taken the following day.

Dosage

Adults over 18 years: 1 tablet up to 3 times a day with water. Leave at least 6 hours between doses. If 1 tablet dose does not control symptoms, a maximum of 2 tablets may be taken up to 3 times a day. Leave at least 6 hours between doses. No more than 6 tablets in 24 hours . To minimise side effects, it is recommended that patients take this medicine with food. Under 18 years: not suitable. For short term-use only. A doctor should be consulted if the symptoms persist or worsen or if the product is required for more than 3 days.

Format and Ingredients

Tablet containing:
Ibuprofen 200mg, Paracetamol 500mg,

Category:

Pack size: 10

Product data © Communications International Group

A division of GROUPE EUROCOM
PAGB
Communications International Group

Communications International Group Ltd 2026,
Linen Hall, 162-168 Regent Street, London W1B 5TB